

# Floodlight Illumination of Infinite Wedges

Matthew Cary<sup>†</sup>      Atri Rudra<sup>†</sup>      Ashish Sabharwal<sup>†</sup>      Erik Vee<sup>§</sup>

<sup>†</sup>Computer Science and Engr, Box 352350, University of Washington, Seattle, WA 98195-2350

<sup>§</sup>IBM Almaden Research Center, Department K53/B2, 650 Harry Road, San Jose, CA 95120

{cary,atri,ashish}@cs.washington.edu      vee@almaden.ibm.com

May 15, 2005

## Abstract

The floodlight illumination problem asks whether there exists a one-to-one placement of  $n$  floodlights illuminating infinite wedges of angles  $\alpha_1, \dots, \alpha_n$  at  $n$  sites  $p_1, \dots, p_n$  in a plane such that a given infinite wedge  $W$  of angle  $\theta$  located at point  $q$  is completely illuminated by the floodlights. We prove that this problem is NP-hard, closing an open problem from 2001 [6]. In fact, we show that the problem is NP-complete even when  $\alpha_i = \alpha$  for all  $1 \leq i \leq n$  (the *uniform* case) and  $\theta = \sum_{i=1}^n \alpha_i$  (the *tight* case). On the positive side, we describe sufficient conditions on the sites of floodlights for which there are efficient algorithms to find an illumination. We discuss various approximate solutions and show that computing any *finite* approximation is NP-hard while  $\varepsilon$ -*angle* approximations can be obtained efficiently.

## 1 Introduction

Illumination problems generalize the well-known art gallery problem (see, e.g., [12, 13]). The task is to mount lights at various sites so that a given region, typically a non-convex polygon, is completely illuminated. The sites can be fixed in advance or not. The region may need to be illuminated from outside (like a soccer field) or from inside (like an indoor gallery). The lights may behave like ideal light bulbs, illuminating all directions equally, or like floodlights, illuminating a certain angle in a certain direction. We use the latter model of floodlights in this paper. This model is quite natural and captures scenarios involving guards or security cameras with restricted angle of vision. Illumination algorithms using floodlights have focused in the past on illuminating the interior of orthogonal polygons [8, 1] and general polygons with restrictions on the floodlights used [2, 9, 7, 16]. There has also been work on the stage illumination problem where one tries to illuminate lines rather than polygons [5].

The problem of illumination of *infinite wedges* by floodlights was introduced by Bose et al. [3]. Refer to Figure 1 for the basic setup and definitions. Given  $n$  sites and  $n$  floodlights, the task is to mount these floodlights, one at each site, and orient them so that a given *generalized wedge* is completely illuminated. Here a generalized wedge refers to an infinite wedge with a continuous finite region adjacent to its apex removed. Formally,

### Definition 1. FLOODLIGHT ILLUMINATION Problem

*Instance:* Sites  $p_1, \dots, p_n$  in  $\mathbb{R}^2$ , angles  $\alpha_1, \dots, \alpha_n > 0$ , and a generalized wedge  $W$  of angle  $\theta$ .

*Question:* Viewing the angles as spans of floodlights, is there an assignment of angles to sites along with angle orientations, that completely illuminates  $W$ ?

A couple of natural restrictions of this problem are the *uniform* case where  $\alpha_i = \alpha$  for all  $1 \leq i \leq n$ , and the *tight* case where  $\sum_{i=1}^n \alpha_i = \theta$ . There is clearly no solution to the problem when  $\sum_{i=1}^n \alpha_i < \theta$ . In general, a solution can be described by a mapping of floodlights to sites along with an angle of orientation for each floodlight. In the tight case, however, every solution can alternatively be described by two permutations  $\sigma$  and  $\tau$  of  $\{1, 2, \dots, n\}$  [14]. Here  $\sigma$  is an ordering of the floodlights and  $p_{\tau(i)}$  is the site at which floodlight of angle  $\alpha_{\sigma(i)}$  is mounted. Floodlight




Figure 1: Basic definitions. W.l.o.g. the axis of  $W$  always points along the negative  $x$ -axis in  $\mathbb{R}^2$ .

orientations in this solution are inferred from  $\sigma$  and  $\tau$  as follows. First  $\alpha_{\sigma(1)}$  is mounted at position  $p_{\tau(1)}$  and oriented so that its upper ray is parallel to the upper boundary  $w_u$  of  $W$ , and for  $2 \leq i \leq n$ ,  $\alpha_{\sigma(i)}$  is mounted at position  $p_{\tau(i)}$  and oriented so that its upper ray is parallel to the lower ray of  $\alpha_{\sigma(i-1)}$ . The variant of the tight floodlight illumination problem where  $\sigma$  is fixed in advance will be called the *restricted case*. When talking about the restricted case, we will think of  $\sigma$  as the identity permutation. Observe that a tight and uniform problem is also effectively restricted because all choices of  $\sigma$  are equivalent. Our results show that in general, for every choice of  $\sigma$ , computing  $\tau$  is NP-complete.

Because of hardness of verification issues surrounding non-algebraic numbers, it is not clear whether the general problem is in the class NP. In fact, it is not obvious that it even has an exponential time solution. Nonetheless, Steiger and Streinu [14] proved that it can indeed be solved in exponential time by formulating it as a bounded quantifier expression in Tarski's algebra [15] and using the result of Grigor'ev [11] on the complexity of deciding the truth value of such expressions. They also proved that the restricted floodlight illumination problem is the dual of a certain *monotone matching* problem with lines and slabs.

The problem in the tight case does not have complications with non-algebraic numbers because the solution, as observed earlier, can be expressed as two permutations on  $\{1, \dots, n\}$ . The tight case of the problem is obviously in NP. However, the exact complexity of this problem has been unknown [6]. We resolve this open question by showing the following.

**Theorem 1.** FLOODLIGHT ILLUMINATION is NP-hard. The tight, restricted, and uniform versions of the problem are NP-complete.

This is an immediate consequence of the discussion of duality in Section 2.2 and our NP-completeness result for a uniform version of monotone matching (Theorem 7). While we know of an NP-hardness reduction from the propositional satisfiability problem 3SAT to the monotone matching problem, the one we give here is from an interesting graph problem, that of finding a *directed disjoint cycle cover*, which we also prove to be NP-complete. Using this latter problem leads to a simplified and more natural proof for the hardness of monotone matching. The problem is also of independent result and a variant of it has subsequently been used by the authors in the context of a finite metric embedding problem [4].

Although the general floodlight illumination problem is NP-hard, many special cases can be solved efficiently. We outline sufficient conditions and list several common site configuration classes for which an efficient greedy algorithm based on duality [14] works correctly in the tight case. There are several natural notions of approximation for the floodlight illumination problem. We consider two of these, a *finite*-approximation where one illuminates all but a finite region of  $W$  and an  $\varepsilon$  *angle*-approximation where one illuminates all but an infinite wedge of small angle  $\varepsilon$  within  $W$ . We prove the following as an immediate consequence of Lemmas 15 and 16.

**Theorem 2.** For the tight floodlight illumination problem, computing a finite-approximation is NP-hard, where as for any  $\varepsilon > 0$ , an  $\varepsilon$  angle-approximation can be constructed in polynomial time.

The rest of the paper is organized as follows. In Section 2 we define the dual monotone matching problem. We also introduce the directed disjoint cycle cover problem and prove it to be NP-complete. In Section 3 we prove the NP-completeness of the floodlight illumination problem by reducing the directed disjoint cycle cover problem to the dual problem of monotone matching. In Section 4 we investigate classes of instances of the tight floodlight illumination problem for which a simple polynomial time algorithm works. Finally, we discuss some notions of approximation for the problem.

## 2 Preliminaries

We begin by defining the monotone matching problem and recapitulating its duality with respect to the restricted floodlight illumination problem. We then define the directed disjoint cycle cover problem and prove its NP-completeness.

### 2.1 Monotone Matching

Suppose we are given  $n$  non-vertical lines in the plane  $\mathbb{R}^2$ ,  $n+1$  vertical lines defining  $n$  finite width vertical slabs, and two points, one on the leftmost vertical line and one on the rightmost. Call this an  $n$ -arrangement of lines, slabs, and points and denote it by  $(L, S, \lambda, \rho)$  where  $L \equiv \{(m_1, c_1), \dots, (m_n, c_n)\}$  is the set of lines  $y = m_i x + c_i$ ,  $1 \leq i \leq n$ ,  $S \equiv \{s_1, \dots, s_{n+1}\}$  is the set of vertical lines  $x = s_j$ ,  $1 \leq j \leq n$ , forming slabs, and  $\lambda$  and  $\rho$  are the two special points on the vertical lines  $x = s_1$  and  $x = s_{n+1}$ , respectively. The reader may want to look at Figures 6-8 for a visual interpretation, although the non-vertical lines in those figures are piecewise linear *pseudolines* defined later. The intersection of any non-vertical line  $\ell$  with a slab  $s$  will be referred to as the *segment* of  $\ell$  spanning  $s$ . A *monotone matching* in  $(L, S, \lambda, \rho)$  is a set of  $n$  line segments, each a portion of a unique line and spanning a unique slab, such that the following holds: (1) the left endpoint of the first segment is above  $\lambda$ , (2) the left endpoint of each subsequent segment is above the right endpoint of the segment in the previous slab, and (3)  $\rho$  is above the right endpoint of the last segment.

**Definition 2.** MONOTONE MATCHING Problem [14]:

*Instance:* An  $n$ -arrangement  $(L, S, \lambda, \rho)$  of lines, slabs, and points in  $\mathbb{R}^2$ .

*Question:* Does this arrangement contain a monotone matching?

Analogous to the floodlight illumination case, define the specialized *uniform* version UNIFORM MONOTONE MATCHING to be the problem where all slabs have the same width. The key technical contribution of this paper is a proof that this variant is NP-complete (see Section 3.2). By the duality argument we discuss shortly, this implies the hardness of the floodlight illumination problem as well, which is our main focus.

### 2.2 Duality Between Floodlight Illumination and Monotone Matching

The restricted floodlight problem can be related to the monotone matching problem through duality [14]. The dual of a point  $p$  with coordinates  $(a, b)$  is the line  $T_p$  with equation  $y = ax + b$ ; the dual of a line  $\ell$  with equation  $y = mx + c$  is the point  $T_\ell$  with coordinates  $(-m, c)$ . It is well known that this dual transformation preserves incidence and height ordering; i.e. if a point  $p$  intersects a line  $\ell$  then their duals  $T_p$  and  $T_\ell$  also intersect, and if  $p$  is above  $\ell$  (w.r.t. the  $y$ -coordinate) then  $T_p$  is above  $T_\ell$ .

We now describe how any restricted floodlight illumination problem can be converted to its dual monotone matching problem  $(L, S, \lambda, \rho)$ . We will use the notation of Figure 1. The duals of the wedge boundaries  $w_l$  and  $w_u$  are the points  $\lambda$  and  $\rho$ ; as  $w_l$  has larger slope in the orientation of the figure, its dual  $\lambda$  has smaller  $x$  coordinate. The points that form the vertical line containing  $\lambda$  are the duals of the lines that are parallel to  $w_l$ . The vertical strip between  $\lambda$  and  $\rho$  corresponds to the wedge angle; the larger the wedge angle, the wider the strip. The line  $q$  that connects  $\lambda$  and  $\rho$  is the dual of the point at which  $w_l$  and  $w_u$  intersect, forming the apex of the wedge. The segment of  $q$  between  $\lambda$  and  $\rho$  corresponds to the lines with slope less than  $w_l$  and greater than  $w_u$  that have common intersection with  $w_l$  and  $w_u$ ; this is exactly the set of lines that lie within  $W \cup W^r$ .

Each site  $p_i$ ,  $1 \leq i \leq n$ , corresponds to a line  $h_i$  which together make up the set of lines  $L$ . As we are in the restricted version of the problem, the angle of the first floodlight can be assumed w.l.o.g. to be  $\alpha_1$ . From our discussion

in Section 1, the tightness of the problem implies that the first floodlight must be oriented so that its upper ray is parallel to  $w_u$ . This floodlight corresponds in the dual to a vertical slab  $S_1$  beginning at  $\rho$  and extending to the left a width proportional to  $\alpha_1$  (if  $S_1$  extends from  $x = s_1$  to  $x = s_2$ , then  $\alpha = \tan^{-1} s_2 - \tan^{-1} s_1$ ). The next floodlight corresponds to a slab vertical  $S_2$  extending to the left of  $S_1$ , continuing to the final floodlight which is a vertical slab  $S_n$  ending at  $\lambda$ .

A solution to the restricted problem is an assignment of sites to floodlights. In the dual this is a 1-1 assignment of lines  $h_i$  to slabs  $S_j$ ,  $1 \leq i, j \leq n$ . The illumination wedge of the first floodlight must overlap  $w_u$ , which corresponds to the right endpoint of the segment of the line  $h_i^1$  assigned to  $s_1$  being above  $\rho$ . Continuing, the right endpoint of the segment associated with  $S_2$  must start above the left endpoint of the segment of  $S_1$ , and so on, until the left endpoint of the segment at  $S_n$  is below  $\lambda$ .

If we flip the dual problem from left to right, in deference to those of us who read from left to right, we see we have reduced the restricted floodlight illumination problem to the monotone matching problem with lines. In the rest of the paper, we will restrict our attention to proving that the monotone matching problem is difficult to solve.

Note that the unrestricted tight illumination problem corresponds to an extended matching problem where the widths of slabs are given and must be arranged in a partition of the slab between  $\lambda$  and  $\rho$  before a matching of segments to slabs is found. The uniform illumination problem corresponds to the uniform matching problem, where the slabs are all of the same width, making, in particular, their order immaterial.

### 2.3 Directed Disjoint Cycle Cover

As a tool for our main result, we prove NP-completeness of an interesting problem on directed graphs which naturally reduces to monotone matching. We also prove that this problem remains NP-complete even when the vertices of the graph are restricted to have small degrees. This latter version is not critical but makes our NP-completeness proof for MONOTONE MATCHING cleaner.

**Definition 3.** DIRECTED DISJOINT CYCLE COVER Problem:

*Instance:* A directed graph  $G = (V, E)$ .

*Question:* Is there a directed disjoint cycle cover for  $G$ , i.e., a set  $\mathcal{C} = \{C_1, \dots, C_k\}$  of vertex disjoint directed cycles in  $G$  such that every vertex  $v \in V$  is in some cycle  $C_i \in \mathcal{C}$ ?

**Theorem 3.** DIRECTED DISJOINT CYCLE COVER is NP-complete, even for graphs with indegree and outdegree each bounded above by 3, as well as for graphs with outdegree exactly 3 and indegree at most 4.

*Proof of Theorem 3.* We begin by giving a proof of the following Lemma that proves NP-completeness of the problem without degree restrictions, and then use it to prove the theorem.

**Lemma 4.** DIRECTED DISJOINT CYCLE COVER is NP-complete.

*Proof.* The problem is clearly in NP because given a description of a set  $\{C_1, \dots, C_k\}$  as a certificate of membership, one can easily verify in polynomial time that each  $C_i$ ,  $1 \leq i \leq k$ , is a directed cycle in the given graph  $G$ , the  $C_i$ 's are vertex disjoint, and they together cover all vertices of  $G$ .

Let VERTEX COVER =  $\{\langle G, k \rangle \mid G \text{ is a directed graph with a vertex cover of size } k\}$ . VERTEX COVER is NP-complete [10]. Our proof of NP-hardness is by a reduction from VERTEX COVER. Our reduction will closely follow the proof of NP-hardness of the HAMILTONIAN CIRCUIT problem given by Garey and Johnson [10]. Let  $\langle G, k \rangle$  be an instance of VERTEX COVER. We will construct a directed graph  $G'$  such that  $G$  has a vertex cover of size  $k$  iff  $G'$  has a directed disjoint cycle cover. In fact,  $G'$  will be such that it has a directed disjoint cycle cover iff it has a directed Hamiltonian circuit. Hence, our proof gives an alternate argument for the NP-hardness of DIRECTED HAMILTONIAN CIRCUIT [10].

The construction can be viewed in terms of component gadgets connected together by communication links. First, the graph  $G'$  has  $k$  "selector" vertices  $a_1, a_2, \dots, a_k$ , which will be used to select  $k$  vertices from the vertex set  $V$  for  $G$ . Second, for each edge in  $e \in E$ ,  $G'$  contains a "cover-testing" component that will be used to ensure that at least one endpoint of  $e$  is among the selected  $k$  vertices. The component for  $e = \{u, v\}$  is illustrated in Figure 2. It has 4 vertices,  $V'_e = \{(u, e, top), (u, e, bot), (v, e, top), (v, e, bot)\}$ , and 6 directed edges,  $E'_e$ , shown in the figure.




Figure 2: Cover-testing component for edge  $e = \{u, v\}$ , along with the three possible configurations of a directed disjoint cycle cover on this component

It is easily seen that any disjoint cycle cover for  $G'$  must involve edges of  $E'_e$  in one of the three configurations shown in Figure 2. This is because any other choice of edges will leave at least one of the gadget vertices impossible to cover. Thus, for example, if the circuit “enters” this component at  $(u, e, top)$ , it must “exit” at  $(u, e, bot)$  and visit either all four vertices in  $V'_e$  or just the two.

Additional edges in our overall construction will serve to join pairs of cover-testing components or to join a cover-testing component to a selector vertex. For each vertex  $v \in V$ , let the edges incident on  $v$  be ordered (arbitrarily) as  $e_{v[1]}, e_{v[2]}, \dots, e_{v[deg(v)]}$ , where  $deg(v)$  denotes the degree of  $v$  in  $G$ . All the cover-testing components corresponding to these edges (having  $v$  as endpoint) are joined together by the following connecting edges:  $E'_v = \{((v, e_{v[i]}, bot), (v, e_{v[i+1]}, top)) : 1 \leq i < deg(v)\}$ . As shown in Figure 3, this creates a single path in  $G'$  that includes exactly the vertices  $(x, y, z)$  having  $x = v$ .




Figure 3: Path joining all cover-testing components for edges from  $E$  having vertex  $v$  as an endpoint

The final connecting edges in  $G'$  join the first and last vertices from each of these paths to every one of the selector vertices,  $a_1, a_2, \dots, a_k$ . These edges are:  $E'' = \{(a_i, (v, e_{v[1]}, top)), ((v, e_{v[deg(v)]}, bot), a_i) : 1 \leq i \leq k, v \in V\}$ . The final graph  $G'$  has  $V = \{a_i : 1 \leq i \leq k\} \cup (\bigcup_{e \in E} V'_e)$  and  $E' = (\bigcup_{e \in E} E'_e) \cup (\bigcup_{v \in V} E'_v) \cup E''$ .  $G'$  can clearly be constructed from  $G$  and  $k$  in polynomial time.

We now argue the correctness of the construction. Suppose first that  $G'$  has a directed disjoint cycle cover  $\mathcal{C}$ . Pick any cycle  $C \in \mathcal{C}$ .  $C$  must traverse vertices of  $G'$  in a particular way, namely, it must start w.l.o.g. at a selector vertex  $a_i$ , go through all cover-testing components corresponding to the edges incident on a particular vertex  $v \in V$ , and either return to  $a_i$  to finish the cycle or return to  $a_j, j \neq i$ , to continue covering in this manner before finally returning to  $a_i$ . This is because (i) each cover-testing component must be covered in one of the three configurations shown in Figure 2, and (ii) edges in  $E'_v$  force  $C$  to traverse all cover-testing components of edges incident on  $v$  in order. Hence, the  $k$  vertices from  $\{a_1, a_2, \dots, a_k\}$  divide  $\mathcal{C}$  into  $k$  paths, each corresponding to a distinct vertex  $v \in V$ . Since  $\mathcal{C}$

must include all vertices from every one of the cover-testing components, and since vertices from the cover-testing component for edge  $e \in E$  can be traversed only by a path corresponding to an endpoint of  $e$ , every edge in  $E$  must have at least one endpoint among those  $k$  selected vertices. Therefore, this set of  $k$  vertices forms the desired vertex cover for  $G$  of size  $k$ .

Conversely, suppose  $\{v_1, v_2, \dots, v_k\} = VC \subseteq V$  is a vertex cover for  $G$  with  $|VC| = k$ . It is readily seen that the following edges form a directed disjoint cycle cover of  $G'$ . From the cover-testing component representing each edge  $e = \{u, v\} \in E$ , choose the edges specified in Figure 2 depending on whether  $\{u, v\} \cap VC$  equals  $\{u\}$ ,  $\{u, v\}$  or  $\{v\}$ . One of these three possibilities must hold because  $VC$  is a vertex cover for  $G$ . Next, choose all the edges in  $E'_{v_i}$  for  $1 \leq i \leq k$ . Finally, choose the edges  $(a_i, (v_i, e_{v_i[1]}, top))$  for  $1 \leq i \leq k$ , edges  $((v_i, e_{v_i[deg(v_i)]}, bot), a_{i+1})$  for  $1 \leq i < k$ , and the edge  $((v_k, e_{v_k[deg(v_k)]}, bot), a_1)$ . This forms a directed Hamiltonian circuit for  $G'$  and, in particular, a directed disjoint cycle cover.  $\square$

We return to the proof of Theorem 3. The restricted degree problem is in NP because, by Lemma 4, the unbounded degree version is in NP. For NP-hardness, suppose we are given an instance  $\langle G, k \rangle$  of VERTEX COVER. Use the construction in the proof of Lemma 4 to obtain a graph  $G'$  which has a directed disjoint cycle cover iff  $G$  has a vertex cover of size  $k$ . Notice that each of the  $k$  selector vertices  $a_i, 1 \leq i \leq k$ , of  $G'$  have indegree as well as outdegree  $n = |V(G)|$  corresponding to the edges in  $E'$ , the  $n$  vertices  $(v, e_{v[1]}, top), v \in V$ , have indegree  $k$  and outdegree 2, and the  $n$  vertices  $(v, e_{v[deg(v)]}, bot), v \in V$ , have indegree 2 and outdegree  $k$ . The rest of the vertices have indegree as well as outdegree 2. We will describe a polynomial time process to convert  $G'$  into a directed graph  $G''$  with indegree and outdegree each bounded above by 3.  $G''$  will have a directed disjoint cycle cover iff  $G'$  does. Further, we will give a very simple way to convert  $G''$  into a graph  $G^*$  that has outdegree exactly 3 and indegree bounded above by 4.  $G^*$  will have a directed disjoint cycle cover iff  $G''$  does. This will finish the proof of the Theorem. A similar construction can alternatively be used to obtain a graph with indegree exactly 3 and outdegree bounded above by 4.

In the rest of the proof, we describe how to reduce the degree of the  $k$  selector vertices. A similar technique can be used to reduce the degrees of the  $2n$  degree- $k$  edge gadget vertices as well; we omit the details of this part for simplicity. The construction of  $G''$  is based on the “degree gadget” shown in Figure 4. Start by making  $G''$  the subgraph of  $G'$  obtained by deleting all edges involving the  $k$  selector vertices  $a_1, \dots, a_k$ . Let  $v_1, v_2, \dots, v_n$  be the vertices of  $G$ . The  $i^{th}$  selector vertex  $a_i$  in  $G'$  has incoming edges from  $(v_j, e_{v_j[deg(v_j)]}, bot)$  and outgoing edges to  $(v_j, e_{v_j[1]}, top)$  for  $1 \leq j \leq n$ . Create vertices  $V_{a_i}^{in} = \{b_{i,j}^{in}, u_{i,j}^{in}, w_{i,j}^{in} \mid 1 \leq j \leq n\}$  and  $V_{a_i}^{out} = \{b_{i,j}^{out}, u_{i,j}^{out}, w_{i,j}^{out} \mid 1 \leq j \leq n\}$  in  $G''$  and connect them as shown in Figure 4. Notice that vertex  $a_i$  in  $G''$  has indegree and outdegree exactly 1, while the other degree gadget vertices have indegree and outdegree between 1 and 3. Overall, the vertices of  $G''$  are  $V(G') \cup \bigcup_{j=1}^k (V_{a_j}^{in} \cup V_{a_j}^{out})$ , all of which have indegree as well as outdegree bounded above by 3.




Figure 4: Degree gadget for the  $i^{th}$  selector vertex

We now argue that  $G''$  has a directed disjoint cycle cover iff  $G'$  does. By construction, cycle covers in the two

graphs behave identically on vertices other than the selector vertices and those of the degree gadgets. The only way to cover  $a_i$  in  $G'$  is to enter from  $(v_x, e_{v_x[\deg(v_x)]}, \text{bot})$  and leave to  $(v_y, e_{v_y[1]}, \text{top})$  for some  $v_x, v_y \in V(G)$ . Similarly, the only way to cover  $a_i$  in  $G''$  is to enter from  $(v_x, e_{v_x[\deg(v_x)]}, \text{bot})$ , traverse through  $b_x^{in}, \dots, b_n^{in}$ , cover  $a_i$ , traverse through  $b_1^{out}, \dots, b_y^{out}$ , and leave to  $(v_y, e_{v_y[1]}, \text{top})$  for some  $v_x, v_y \in V(G)$ . The remaining degree gadget vertices in  $V_{a_i}^{in} \cup V_{a_i}^{out}$  must be covered in 3-cycles  $(b_{i,j}^x, w_{i,j}^x, w_{i,j}^x)$  or 2-cycles  $(u_{i,j}^x, w_{i,j}^x)$ . This shows that cycle covers in  $G'$  and  $G''$  behave essentially identically even on the selector vertices and the degree gadgets. Hence,  $G'$  has a directed disjoint cycle cover iff  $G''$  has one.

$G^*$  is constructed from  $G''$  as follows. Start with  $G^* = G''$ . Let  $K_4$  denote the directed complete graph on 4 vertices. Note that each vertex of  $K_4$  has indegree as well as outdegree exactly 3. Let  $p = |V(G'')|$  and  $q$  be the sum of the outdegrees of the vertices of  $G''$ . Create  $3p - q$  copies of  $K_4$  and add them to  $G^*$ . For each outdegree 2 vertex  $v$  in  $G$ , add an outgoing edge to an arbitrarily selected vertex of an unmarked copy  $K_4$  and mark that copy (see Figure 5). Similarly, do this for each outdegree 1 vertex, connecting it to two copies of  $K_4$ . The number of edges added thus is  $3p - q$ , which exactly matches the number of copies of  $K_4$  created in  $G^*$ . Every vertex of  $G^*$  now has outdegree exactly 3. The indegree of the original vertices from  $G''$  hasn't changed and is therefore at most 3. The indegree of exactly one vertex in each copy of  $K_4$  has changed, and it has increased to 4. Hence,  $G^*$  has the desired degree properties.




Figure 5: Using copies of  $K_4$  to make outdegrees exactly 3

Since the copies of  $K_4$  do not have any outgoing edge, the only cycles they can be involved in a disjoint cycle cover stay within that copy. The new edges added between  $G''$  and the copies of  $K_4$  can never be used in a cycle. This ensures that if  $G^*$  has a disjoint cycle cover, then  $G''$  does too. On the other hand, if  $G''$  has a disjoint cycle cover, then one for  $G^*$  consists of cycles corresponding to the cover in  $G''$  along with one 4-cycle for each copy of the  $K_4$ . Hence,  $G^*$  has a directed disjoint cycle cover iff  $G''$  does.  $\square$

### 3 NP-Completeness of Monotone Matching

We first begin by proving the NP-Completeness of a generalization of the monotone matching problem in Section 3.1. The proof for the NP-Hardness is easier to present and would convey all the main ideas in our reduction from the directed disjoint cycle cover. We then prove the NP-Hardness of the monotone matching problem in Section 3.2.

#### 3.1 Monotone Matching with Pseudolines

We begin by defining a variant of the monotone matching problem that uses *pseudolines* instead of lines. A pseudoline is a curve in  $\mathbb{R}^2$  that intersects any vertical line in exactly one point. A collection of pseudolines is a set of pseudolines no two of which intersect more than once. For computational purposes, we shall assume that the point of intersection of two pseudolines can be computed efficiently from their input representation.  $n$ -arrangements of pseudolines, slabs, and points, and monotone matchings with pseudolines are defined analogous to the case of straight lines. The problem PSEUDOLINE MONOTONE MATCHING is also defined analogously. The pseudolines used in our constructions will all be piecewise linear functions. In this section, we prove the following result.

**Lemma 5.** PSEUDOLINE MONOTONE MATCHING is NP-complete.

We begin by defining some configurations, or gadgets, of pseudolines that will be useful for our proof. Our construction for the straight line matching problem will follow by reconstructing these gadgets from straight lines.

In our reduction we would like to make sure that only a subset of the pseudolines can be used in some particular slabs. This is achieved by perhaps the most important gadget in our reduction which we call the *forcing gadget* (see Figure 6). This is a sequence of slabs associated with pseudolines that forces the pseudoline used previous to the gadget to end below a chosen point, and the pseudoline used after the gadget to start above another chosen point. Let  $R$  denote the *region of interest* for the monotone matching problem, i.e., a rectangular region covering horizontally all the slabs and covering vertically all intersections of pseudolines in  $PL$  with these slabs and the special points  $\lambda$  and  $\rho$ . The *sky* region begins above  $R$  and the *ground* region begins below  $R$ . Each of these regions is further divided into *high* and *low* regions. The idea is that once a pseudoline reaches high sky, it cannot be used anymore because no pseudoline comes down from high sky to  $R$ . Similarly, no pseudoline can be used until it leaves low ground because no pseudoline comes down from  $R$  to low ground. The low sky and high ground regions are intermediate regions where pseudolines of negative slope that are part of the forcing gadget reside before and after they are called upon for use in the gadget.

Every forcing gadget  $F$  is associated with two pseudolines  $h$  and  $\ell$  (see Figure 6). We call these *auxiliary* pseudolines.  $h$  is parallel to the boundary between low sky and  $R$  till it goes down before its corresponding slab  $A$  and finally it is parallel to the boundary between high ground and  $R$ . The situation for the other pseudoline is complementary, that is,  $\ell$  is parallel to the boundary between low ground and high ground till it reaches its slab  $A$  and is parallel to the boundary between low and high sky after its corresponding slab  $B$ . For any pair of gadgets  $F$  and  $F'$  (with pseudolines  $h'$  and  $\ell'$ ) such that  $F$  appears before  $F'$ ,  $h$  is below  $h'$  in the low sky region and is below  $h'$  in the high ground region while  $\ell$  is above  $\ell'$  in the low ground region and above  $\ell'$  in the high sky region. Note that this implies that  $h$  and  $h'$  do not intersect and neither do  $\ell$  and  $\ell'$ . Further we require that the point  $s$  in  $F$  is below  $h'$ . It is easy to see that given the positions of the forcing gadgets their corresponding pseudolines and the high/low sky and ground regions can be constructed in polynomial time.




Figure 6: The forcing gadget. The arrows show how any lines used before or after the gadget are constrained.

**Lemma 6 (Forcing Gadget Property).** *Refer to Figure 6 for the labeling of forcing gadgets. Given a monotone matching instance, for all pairs  $F$  and  $F'$  of forcing gadgets composed of  $h, \ell, s$  and  $h', \ell', s'$ , respectively, in any valid matching,  $h$  is used in slab  $A$  and  $\ell$  is used in slab  $B$ . This holds for any number of additional lines, as long as they appear above the ground and below the sky.*

*Proof.* Recall from our construction of the forcing gadgets and their corresponding pseudolines,  $h$  and  $h'$ , do not intersect, and the point  $s$  of  $F$  is below  $h'$  if  $F'$  follows  $F$ .

$\ell$  cannot be used before slab  $B$ , as it begins in the ground, no other lines intersect  $\ell$  before  $B$ , and the ground is below the starting point. If  $\ell$  is used after slab  $B$ , then the line used subsequently must begin in the high sky. As the only lines in the high sky are from previous forcing gadgets, and there are no intersections in the high sky, then all future lines chosen will be in the high sky. As the high sky is above  $\rho$ , any such matching must end above  $\rho$ , and so will not be a valid matching.

Now, given that  $\ell$  must be used in slab  $B$ , as its starting point  $s$  is in high ground, the only precursors it could have is  $h$ , some  $h'$  from a forcing gadget before  $F$ , or some line from the low ground. As argued in the beginning of this proof, no lines in the low ground can be used. We prove that the precursor could not come from an  $h'$  preceding  $F$  by induction on the number of forcing gadgets. If  $F$  is the first forcing gadget, then  $h$  must be used; if there are forcing gadgets before  $F$ , then we can inductively assume that all the corresponding  $h'$  were used in their corresponding gadgets, so that  $h$  must be used in slab  $A$ .

Finally, as our arguments were based on the presence or absence of pseudolines in the ground and sky, any number of additional lines appearing between the ground and sky do not affect the behavior of the forcing gadget.  $\square$

*Proof of Lemma 5.* This problem is in NP because a potential monotone matching can be easily verified in polynomial time. The proof of NP-hardness is by a reduction from DIRECTED DISJOINT CYCLE COVER, which, by Theorem 3, is NP-complete. Suppose we are given a directed graph  $G$ . We will construct an arrangement  $(PL, S, \lambda, \rho)$  of pseudolines, slabs, and points such that  $G$  has a disjoint cycle cover iff  $(PL, S, \lambda, \rho) \in$  PSEUDOLINE MONOTONE MATCHING. By Theorem 3, we may assume the outdegree of all vertices in  $G$  is exactly 3 and the indegree is at most 4. W.l.o.g. we will assume that every vertex in  $G$  has indegree at least one, for if not then  $G$  does not have a disjoint cycle cover and can be easily mapped to a trivial instance of monotone matching with no solution.

We need two types of graph-related gadgets. Let  $G = (V, E)$ . We will have gadgets  $\mathbf{In}(v)$  and  $\mathbf{Out}(u)$  for  $u, v \in V$  as shown in Figure 7. Let  $\mathcal{I}(v) \subset E$  be in the in-edges of  $v$ , and let  $\mathcal{O}(u) \subset E$  be the out-edges of  $u$ . By our choice of  $G$ ,  $|\mathcal{I}(v)| \leq 4$  and  $|\mathcal{O}(u)| = 3$ . The gadgets  $\mathbf{In}(v)$  and  $\mathbf{Out}(u)$  will allow us to select exactly one predecessor of  $v$  and exactly one successor of  $u$ , respectively, leading to a directed disjoint cycle cover as long as the predecessors and successors are consistent. Selecting a successor will be done indirectly by selecting and ruling out exactly  $|\mathcal{O}(u)| - 1$  out-edges of  $u$ .




Figure 7: Graph gadgets  $\mathbf{In}(v)$  and  $\mathbf{Out}(u)$  are shown in (a) and (b), respectively. The arrows denote sites set by forcing gadgets placed between the graph gadgets as shown in (c).  $|\mathcal{I}(v)| \leq 4$  and  $|\mathcal{O}(u)| = 3$ .

Let  $n = |V|$  and  $m = |E|$ . We will use  $m$  primary pseudolines, each corresponding to an edge in  $E$ . We will abuse notation and talk of an edge  $e \in E$  as being used in a particular slab; this will mean that the primary pseudoline corresponding to  $e$  is used at that slab. There will be a number of auxiliary pseudolines used in forcing gadgets. The  $\mathbf{Out}(\cdot)$  gadgets and  $\mathbf{In}(\cdot)$  gadgets will be arranged in sequence as shown in Figure 8. The primary pseudoline corresponding to edge  $(u, v)$  will first pass through  $\mathbf{Out}(u)$ , and then pass through  $\mathbf{In}(v)$ . Each pseudoline will intersect another pseudoline at most once, depending on the relationship between their  $\mathbf{In}(\cdot)$  and  $\mathbf{Out}(\cdot)$  gadgets.

We claim that when arranged as in Figure 8 along with appropriate forcing gadgets for each  $\mathbf{In}(\cdot)$  and  $\mathbf{Out}(\cdot)$  gadget, exactly one  $e \in \mathcal{I}(v_i)$  is used in  $\mathbf{In}(v_i)$  and exactly one  $e \in \mathcal{O}(v_i)$  is not used in  $\mathbf{Out}(v_i)$ , for  $1 \leq i \leq n$ . To see this, refer to Figure 7. Because of the forcing gadgets, exactly one pseudoline must be used in  $\mathbf{In}(v_i)$  and must correspond to an edge  $e \in \mathcal{I}(v_i)$ . Similarly, exactly two pseudolines must be used in  $\mathbf{Out}(v_i)$ , leaving out exactly




Figure 8: Overall view of the reduction from DIRECTED DISJOINT CYCLE COVER. Forcing gadgets appear between  $\mathbf{In}(\cdot)$  and  $\mathbf{Out}(\cdot)$  gadgets, to match the arrows in Figure 7, except for  $\mathbf{Out}(v_1)$  and  $\mathbf{In}(v_n)$ , where the first and last arrows, respectively, determine the starting and ending points  $\lambda$  and  $\rho$  for the monotone matching instance.

one pseudoline corresponding to an edge  $e \in \mathcal{O}(v_i)$ . Furthermore, the edges selected in this way are consistent, i.e., if edge  $(u, v)$  is selected in  $\mathbf{In}(v)$ , then it is also the only edge left unselected in  $\mathbf{Out}(u)$ .

A directed disjoint cycle cover of  $G$  is equivalent to a permutation  $\pi$  on the vertices, where  $\pi(v)$  is the predecessor of  $v$  in the cycle containing  $v$ . If such a permutation exists then a monotone matching exists: don't select the edge corresponding to  $\pi^{-1}(u)$  in  $\mathbf{Out}(v)$  and select the edge corresponding to  $\pi(v)$  in  $\mathbf{In}(v)$ . Conversely, if a monotone matching exists then the permutation  $\pi$  can be recovered by letting  $\pi(v)$  correspond to the edge that is used in  $\mathbf{In}(v)$ . This completes the reduction.  $\square$

### 3.2 Uniform Monotone Matching

We now prove NP-completeness of the uniform case with straight lines. As mentioned earlier, NP-hardness of this problem subsumes NP-hardness of PSEUDOLINE MONOTONE MATCHING proved in the previous section. The argument, however, is more involved and reuses many key concepts developed in the earlier proof.

**Theorem 7.** UNIFORM MONOTONE MATCHING is NP-complete.

*Proof.* Being a sub-problem of PSEUDOLINE MONOTONE MATCHING, the problem is in NP. We prove NP-hardness by a reduction from DIRECTED DISJOINT CYCLE COVER in a manner similar to the proof of Lemma 5. Let  $G = (V, E)$  be a graph of outdegree exactly 3 and indegree at most 4. We will construct an arrangement  $(L, S, \lambda, \rho)$  of lines, slabs, and points such that  $G$  has a directed disjoint cycle cover iff  $(L, S, \lambda, \rho) \in$  UNIFORM MONOTONE MATCHING.

The key changes from the previous proof are the new construction of the forcing gadgets and the addition of a “buffer” zone between the  $\mathbf{In}$  and  $\mathbf{Out}$  gadgets to convert primary pseudolines into straight lines. The basic idea of the buffer zone is to make the lines behave as parallel (i.e. non-intersecting) lines in the region of interest as required in the graph gadgets of Figure 7. We first spell out the details of construction of the lines corresponding to the edges of  $G$  and then specify how to construct the forcing gadgets with lines. For the rest of the proof, we will assume w.l.o.g. that the slab width  $w = 1$ .

The graph gadgets are arranged as shown in Figure 9 with a buffer area consisting of  $b - 4n$  slabs (or  $\frac{b-4n}{2}$  forcing gadgets). We will use a coordinate system as shown in Figure 9 for the ease of exposition. The graph gadget  $\mathbf{In}(v_j)$  has its forced start point at  $(b + 3j - 2, n - j)$  and forced end point at  $(b + 3j - 1, n - j + 1)$  while the graph gadget  $\mathbf{Out}(v_i)$  has its forced start and end points at  $(4i - 3, -i)$  and  $(4i - 1, 1 - i)$ , respectively. The line corresponding

to edge  $(v_i, v_j)$  connects the points  $(-1, -i)$  and  $(b + 3j - 3, n - j)$ . Note that for any two consecutive  $\mathbf{In}(\cdot)$  and  $\mathbf{Out}(\cdot)$  gadgets, there is space for two slabs which would be used up by two forcing gadgets.




Figure 9: Overall view of the reduction from DIRECTED DISJOINT CYCLE COVER for the uniform case. Forcing gadgets appear in the buffer area and between  $\mathbf{In}(\cdot)$  and  $\mathbf{Out}(\cdot)$  gadgets to match the arrows in Figure 7, except for  $\mathbf{Out}(v_1)$  and  $\mathbf{In}(v_n)$  where the first and last arrows, respectively, determine the starting and ending points  $\lambda$  and  $\rho$  for the uniform monotone matching instance. For clarity, only the lines corresponding to outgoing edges from  $v_1$  are shown: the edges are  $(v_1, v_1)$ ,  $(v_1, v_2)$  and  $(v_1, v_n)$ .

Fix  $b = 10n^2$ . We now show that the lines corresponding to the outgoing edges from any graph gadget do not intersect each other in the horizontal intervals  $[0, 4n]$  and  $[b, b + 3n]$ , that is, they are effectively parallel as needed in Figure 7. For any edge  $(v_i, v_j)$  the slope of the corresponding line is bounded from above by  $m = \frac{n - (-n)}{b} = \frac{2n}{b}$ . The maximum y coordinate for the line corresponding to  $(v_i, v_j)$  in the horizontal interval  $[0, 4n]$  is given by  $y_m = -i + m \cdot 4n$ . Substituting the value of  $m$  and  $b$ , we have  $y_m < -i + 1$ . Thus, no two lines intersect in the horizontal interval  $[0, 4n]$ . One can similarly show that no two lines intersect in the horizontal interval  $[b, b + 3n]$ .

To complete the proof we specify the forcing gadgets. We need  $\frac{b}{2}$  forcing gadgets ( $n$  each from the  $\mathbf{In}$  and  $\mathbf{Out}$  gadgets and  $\frac{b-4n}{2}$  from the buffer area). The scheme is presented in Figure 10. The high ground, low ground, low sky, and high sky regions are  $-n^4 \leq y < -n$ ,  $y > -n^4$ ,  $n < y \leq n^4$ , and  $y > n^4$ , respectively. For the  $k^{\text{th}}$  forcing gadget, the  $h$  line connects  $p_0$ , which is at height  $n^4 - 2kn$ , and  $p_1$ , the forced end point at the boundary of slab  $A$ . The  $\ell$  line of the gadget connects  $q_0$  (which is one higher than  $p_2$ , the intersection of  $h$  with the boundary of slab  $B$ ) and  $q_1$ , the forced start point at the boundary of slab  $B$ .

We claim that the height of  $q_2$  is (strictly) bounded from above and below by  $n^4 - 2kn$  and  $n^4 - 2(k + 1)n$ , respectively. By our overall construction in Figure 9,  $p_1$  is always higher than  $q_1$  and their difference is bounded from above by  $n$ . The bounds on the height of  $q_2$  follow from recalling that each slab has width one.

We now argue that in a valid uniform monotone matching,  $h$  is chosen in slab  $A$  and line  $\ell$  is chosen in slab  $B$  as was proved in Lemma 6. We first show that  $\ell$  is selected in slab  $B$ . For the sake of contradiction assume that  $\ell$  was chosen after slab  $B$ . By the lower bound on the height of  $q_2$ , we cannot come down from the sky as all the  $h'$  lines of the subsequent gadgets are below  $q_2$  which is a contradiction as the ending point is below the sky. If  $\ell$  were chosen before slab  $B$ , its precursor cannot include any  $\ell'$  line from a previous forcing gadget because then the starting point  $\lambda$  would have to be below ground, which is not the case. It also cannot have any  $h'$  line from a previous forcing gadget as a precursor as all those lines are higher than  $\ell$  by construction. Hence,  $\ell$  is selected in slab  $B$ . From the above argument the only precursors of  $\ell$  could be  $h$  or some  $h'$  from some previous forcing gadget which, as before by induction, is impossible. Thus,  $h$  is selected in slab  $A$  and  $\ell$  in slab  $B$ , fulfilling the requirements of the forcing gadget.  $\square$




Figure 10: Forcing gadget constructed with straight lines.

**Corollary 8.** MONOTONE MATCHING is NP-complete.

*Proof.* Membership in NP follows from an argument identical to the one for UNIFORM MONOTONE MATCHING. NP-hardness follows from Theorem 7 because UNIFORM MONOTONE MATCHING trivially reduces to this problem.  $\square$

## 4 Illumination Algorithms for the Tight Case

In this section we look at algorithms to solve the floodlight illumination problem in the tight case. We characterize several special cases that can be solved in polynomial time. We also give approximation algorithms for the problem. For this section, we will use the notations and definitions from Section 1. We begin with some properties of floodlight illuminations which will be used later in this section but may also be of independent interest.

### 4.1 Properties of Floodlight Illumination

The first lemma shows how the position of certain floodlights are fixed by the problem instance. Next we prove a necessary condition for the existence of a solution. We then state a lemma which will be used in proving hardness of certain kind of approximations to the floodlight illumination problem. We end with a brief mention of a variant of the problem that is easy to solve.

**Lemma 9.** *In the tight case, any floodlight illuminating at infinity the upper boundary  $w_u$  of a generalized wedge  $W$  must be parallel to and located above  $w_u$ .*

*Proof.* Let the floodlight  $f$  illuminating  $w_u$  be mounted at site  $p_u$ . As we are considering the tight case, the upper boundary  $f_u$  of the region illuminated by  $f$  must be parallel to  $w_u$  (see discussion in Section 1. If  $p_u$  is below  $w_u$ ,  $f$  will not illuminate an infinite slice  $S$  of  $W$  including the boundary  $w_u$ .  $\square$

**Lemma 10.** *A tight floodlight illumination problem on generalized wedges has a solution only if there is at least one site in the reverse wedge.*

*Proof.* Suppose there are no sites in the reverse wedge. We will derive a contradiction by arguing about the dual of the problem as defined in Section 2.2. The lines in the dual corresponding to sites fall into three categories: those above  $\rho$  and  $\lambda$  (the duals of  $w_u$  and  $w_\ell$ ), corresponding to sites above the wedges; those below the duals  $\rho$  and  $\lambda$ , corresponding to sites below the wedge; and those that run between  $\rho$  and  $\lambda$ , corresponding to sites in the forward wedge (recall that we have assumed that there are no sites in the reverse wedge). As we are in the tight case, if there are sites in the forward wedge, we cannot possibly have a solution. Hence all points are either above or below both of the dual starting and end points. In particular, no line above both the dual points intersects a line below both the




Figure 11: Two cases for the shape of overall illuminated region in the tight case

dual points in the problem slab defined by the dual points. This means that we cannot use any of the lines below the starting point and hence, we cannot have a matching as there are not enough lines which can be used.  $\square$

**Lemma 11.** *Suppose the sum of the angles of  $n$  floodlights is  $\alpha$ . If they illuminate a wedge  $W$  of angle  $\alpha$ , then the overall illuminated region is of the form  $W' \cup S$ , where  $W' \supseteq W$  is a wedge of angle  $\alpha$  aligned with  $W$  and  $S$  is a finite region.*

*Proof.* We prove by induction on  $n$  the weaker statement that ignores the requirement  $W' \supseteq W$  above. This will, however, suffice for the proof because if  $W'$  did not include all of  $W$ ,  $W' \cup S$  would also not include all of  $W$  since  $S$  is finite. This will contradict the precondition.

For the base case of  $n = 1$ ,  $W' = W$  and  $S = \phi$ . For  $n > 1$ , note that the floodlight angles are tight relative to  $W$ . Hence, the first  $n - 1$  floodlights must together cover some wedge of angle  $\alpha - \alpha_n$ , where  $\alpha_n$  is the angle of the  $n^{\text{th}}$  floodlight (see Figure 11). By induction, the region illuminated by the first  $n - 1$  floodlights is of the form  $W'' \cup S'$ , where  $W''$  is a wedge of angle  $\alpha - \alpha_n$  and  $S'$  is finite. Again, since the angles are tight, the only way to extend this region to cover  $W$  of angle  $\alpha$  is to mount the  $n^{\text{th}}$  floodlight  $f$  at a site  $p$  that is above the lower boundary  $w_l$  of the already illuminated region and have its upper boundary  $f_u$  aligned with  $w_l$ . Let  $W'$  be the wedge of angle  $\alpha$  defined by the upper boundary  $w_u$  of the already illuminated region and the lower boundary  $f_l$  of  $f$ . As seen from the two cases in Figure 11, the overall illuminated region is  $W' \cup S$ , where  $S$  is finite.  $\square$

Finally, we mention a relaxation of the problem which makes it easy. Two **movable sites** can always solve any tight problem instance: assign two arbitrarily chosen first and last floodlights (the ones parallel to the wedge boundaries) to the movable sites, and move these sites back and inside the reverse wedge far enough so that every other site is within the reverse of the residual wedge. Now use Fact 12.

## 4.2 A Greedy Algorithm

We briefly describe a duality-based greedy algorithm  $\mathcal{A}_{\text{greedy}}$  given by Steiger and Streinu [14] for the floodlight illumination problem which takes an additional input: the order in which the floodlight angles are chosen, that is, permutation  $\sigma$  from Section 1. Note that for the uniform case, where each floodlight angle is the same, the permutation  $\sigma$  does not come into play and  $\mathcal{A}_{\text{greedy}}$  is applicable. At each step,  $\mathcal{A}_{\text{greedy}}$  assigns the current floodlight angle in  $\sigma$  to the position  $p$  which would leave the maximum number of positions inside the reverse wedge of the residual wedge obtained by placing the current floodlight angle on  $p$ . There is also a natural interpretation of  $\mathcal{A}_{\text{greedy}}$  in the dual monotone matching problem, where one chooses a line for a slab that maximizes the number of choices for the next slab. We will refer to this corresponding dual monotone matching algorithm as  $\mathcal{A}_{\text{greedy}}^{MM}$ . We complete the description of  $\mathcal{A}_{\text{greedy}}$  by stating a simple property of it.

**Fact 12.** *If all sites are contained inside the reverse wedge  $W^r$ , then  $\mathcal{A}_{\text{greedy}}$  successfully illuminates  $W$  after any assignment of floodlights to positions. Equivalently, if the first special point in a monotone matching problem is below all lines and the second is above all, then  $\mathcal{A}_{\text{greedy}}^{MM}$  successfully finds a matching.*

### 4.3 Special Site Configurations

In this section we consider special illumination problems where the sites are restricted to obey certain properties. This allows us to characterize cases where  $\mathcal{A}_{\text{greedy}}^{MM}$  produces the right answer for the corresponding tight floodlight illumination problem.

**Definition 4.** Sites  $p_1, p_2, \dots, p_n$  are *angle-separated* with respect to wedge  $W$  of angle  $\alpha$  if  $p_i \notin W_j$  for every  $1 \leq i \neq j \leq n$ , where  $W_j$  is the wedge of angle  $\alpha$  located at  $p_j$  and aligned with  $W$  (see Figure 12). Sites in a floodlight illumination problem are *angle-separated* if they are angle-separated with respect to the wedge to be illuminated.

We note that an equivalent way of defining angle-separation with respect to a wedge  $W$  with boundary slopes  $m_u$  and  $m_l$  is to require that for all site pairs  $(p, p')$ , the line joining  $p$  and  $p'$  has slope not in  $[m_l, m_u]$ . While the former definition is natural for the proof of the following lemma, this latter definition might be more convenient for algorithmic implementations.




Figure 12: Angle separation implies all sites must be in  $W^r$  for a solution to exist.

**Lemma 13.** *If the sites in a tight floodlight illumination problem on generalized wedges are angle-separated, there is no solution unless all sites are contained in the reverse wedge.*

*Proof.* Consider the dual (refer to Section 2.2). The wedge  $W_j$  for site  $p_j$  corresponds in the dual to the segment of the line  $\ell_{p_j}$  between the starting and ending points, that is, the problem slab. If there is a site that is not in the reverse wedge then there exists a line above the endpoint or a line below the starting point. This implies that one cannot use at least one line in the matching and thus, a matching cannot exist.  $\square$

**Proposition 14 (Sufficient Condition for  $\mathcal{A}_{\text{greedy}}$ ).** *If the sites in a tight floodlight illumination problem on generalized wedges are angle-separated, then  $\mathcal{A}_{\text{greedy}}$  always produces the right answer.*

*Proof.* If all sites are contained inside the reverse wedge, by Fact 12, a solution is always found by  $\mathcal{A}_{\text{greedy}}$  for any assignment of floodlights to sites. On the other hand, if at least one site is outside the reverse wedge, by Fact 13, there is no solution and  $\mathcal{A}_{\text{greedy}}$ , of course, doesn't find one.  $\square$

It follows that the floodlight illumination problem for generalized wedge  $W$  of angle  $\alpha$  and with boundaries of slopes  $m_u$  and  $m_l$  is easy to solve when, for instance, all sites are on a straight line whose slope is not in  $[m_l, m_u]$ , or on a circular arc whose endpoint tangents have slope not in  $[m_l, m_u]$ , or in a vertical zig-zag pattern with angle greater than  $\alpha$ , etc. (see Figure 13).




Figure 13: A few natural site configurations for which the problem is easy.

#### 4.4 Approximate Illuminations

Theorem 1 motivates the study of approximation algorithms for the floodlight illumination problem. There are several natural ways to approximate wedge illumination. After giving precise definitions of some of these, we prove a negative result that illuminating all but a finite portion of a wedge is, in the tight case, not any easier than illuminating the whole wedge. This motivates other reasonable notions of approximation that leave unlit relatively small but infinite regions of the wedge.

**Definition 5.** Let  $\mathcal{F}$  be an illumination of a wedge  $W$ .

1.  $\mathcal{F}$  is a *finite-approximation* if it illuminates  $W \setminus S$ , where  $S$  is a finite region.
2.  $\mathcal{F}$  is an  $\varepsilon$  *angle-approximation* if it illuminates  $W \setminus S_\varepsilon$ , where  $S_\varepsilon$  is a union of wedges whose total angle is at most  $\varepsilon$ .

**Lemma 15.** *There is a solution to a tight floodlight illumination problem on a wedge  $W$  if and only if there is a finite-approximation to it.*

*Proof.* We prove the sufficient condition. Suppose there is a finite-approximate illumination  $\mathcal{F}$  for  $W$ . Let  $W$  be of angle  $\alpha$ . By definition,  $\mathcal{F}$  must illuminate a wedge  $W^*$  of angle  $\alpha$  that is aligned with  $W$  but is possibly contained strictly within  $W$ . Since the floodlight angles are tight relative to  $W^*$ , by Lemma 11, the overall region illuminated by  $\mathcal{F}$  is of the form  $W' \cup S$ , where  $W'$  is a wedge of angle  $\alpha$  aligned with  $W^*$  (and hence with  $W$ ) and  $S$  is finite. If  $W \not\subseteq W'$ , then  $W' \setminus W$  is an infinite region  $R$ . As  $S$  is finite,  $W' \cup S$  will not cover an infinite portion of this infinite region  $R$  of  $W$ , contradicting the fact that  $\mathcal{F}$  illuminates all but a finite region of  $W$ . It follows that  $W \subseteq W'$ , implying that  $W$  is completely illuminated by  $\mathcal{F}$  and providing an exact solution. The other direction of the proof is trivial.  $\square$

This Lemma implies that computing a finite-approximation is NP-hard because computing the exact solution is. It also implies that there is a solution to the tight floodlight problem on a generalized wedge  $W$  iff there is a solution to the tight floodlight problem on the underlying normal wedge  $W'$ . In this sense, generalized wedges don't make the problem any harder. However, they provide a convenient tool for analysis, allowing, for instance, stronger inductive claims.

Note that  $\varepsilon$  angle-approximate illumination only requires all but  $\varepsilon$  of the wedge is illuminated “at infinity”. It would be interesting to find an algorithm for the stronger approximation where the resulting illuminated area is a smaller wedge but located at the same apex as  $W$ .

**Lemma 16.** *For any  $\varepsilon > 0$ , an  $\varepsilon$  angle-approximation to the tight floodlight problem can be found efficiently.*

*Proof.* An  $\varepsilon$  angle-approximation can be achieved by adding two movable sites  $p_a$  and  $p_b$ , adding two floodlights  $f_a$  and  $f_b$  of angle  $\varepsilon/2$  each, reducing any one original floodlight angle by  $\varepsilon$ , and proceeding as follows. Mount floodlight  $f_a$  at site  $p_a$ , orient it so that its upper boundary is parallel to and illuminates the upper boundary  $w_u$  of  $W$ , and move it far and low enough in  $W^r$  so that all other sites are above its lower boundary. Perform a similar operation on  $f_b$

and  $p_b$  starting with the lower boundary  $w_l$  of  $W$ . The region  $W'$  of  $W$  not illuminated by these two floodlights is a generalized wedge of angle  $\alpha - \varepsilon$ , where  $\alpha$  is the wedge angle of  $W$ . Further, all remaining sites live within  $W''$ . By Lemma 12, we can illuminate  $W''$  exactly using the remaining floodlights. Now remove  $f_a$  and  $f_b$ , and add angle  $\varepsilon$  back to the floodlight whose angle was reduced. This completes the  $\varepsilon$  angle-approximation.  $\square$

## Acknowledgments

The authors would like thank Neva Cherniavsky for introducing the problem, and Gidon Shavit and William Pentney for helpful discussions.

## References

- [1] J. Abello, V. Estivill-Castro, T. C. Shermer, and J. Urrutia. Illumination of orthogonal polygons with orthogonal floodlights. *International Journal of Computational Geometry and Applications*, 8(1):25–38, 1998.
- [2] J. Bagga, L. Gewali, and D. Glasser. The complexity of illumination polygons by  $\alpha$ -flood-lights. In *Proceedings of the 8th Canadian Conference on Computational Geometry*, pages 337–342, Ottawa, Canada, Aug. 1996.
- [3] P. Bose, L. Guibas, A. Lubiw, M. Overmars, D. Souvaine, and J. Urrutia. The floodlight problem. *International Journal of Computational Geometry and Applications*, 7(1/2):153–163, 1997.
- [4] M. Cary, A. Rudra, and A. Sabharwal. On the hardness of embeddings between two finite metrics. In *International Colloquium on Automata, Languages and Programming*, Lisboa, Portugal, July 2005. To appear.
- [5] J. Czyzowicz, E. Rivera-Campo, and J. Urrutia. Optimal floodlight illumination of stages. In *Proceedings of the 5th Canadian Conference on Computational Geometry*, pages 393–398, Waterloo, Canada, Aug. 1993.
- [6] E. Demaine and J. O’Rourke. Open problems from CCCG 2001. In *Proceedings of the 13th Canadian Conference on Computational Geometry*, Waterloo, Canada, Aug. 2001.
- [7] V. Estivill-Castro, J. O’Rourke, J. Urrutia, and D. Xu. Illumination of polygons with vertex lights. *Information Processing Letters*, 56(1):9–13, 1995.
- [8] V. Estivill-Castro and J. Urrutia. Optimal floodlight illumination of orthogonal art galleries. In *Proceedings of the 6th Canadian Conference on Computational Geometry*, pages 81–86, Saskatchewan, Canada, Aug. 1994.
- [9] V. Estivill-Castro and J. Urrutia. Two-floodlight illumination of convex polygons. In *Proceedings of the 4th International Workshop on Algorithms and Data Structures*, volume 955 of LNCS, pages 62–73, Kingston, Canada, Aug. 1995. Springer Verlag.
- [10] M. R. Garey and D. S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman and Company, 1979.
- [11] D. Y. Grigor’ev. Complexity of deciding Tarski algebra. *Journal of Symbolic Computation*, 5:65–108, 1998.
- [12] J. O’Rourke. *Art Gallery Theorems and Algorithms*. Oxford University Press, 1987.
- [13] T. C. Shermer. Recent results in art galleries. *Proceedings of the IEEE*, 80(9):1384–1399, 1992.
- [14] W. L. Steiger and I. Streinu. Illumination by floodlights. *Computational Geometry*, 10:57–70, 1998.
- [15] A. Tarski. *A Decision Method for Elementary Algebra and Geometry*. University of California Press, 1948.
- [16] C. D. Tóth. Illuminating polygons with vertex  $\pi$ -floodlights. In *Proceedings, Part I, of the International Conference on Computational Science (ICCS)*, volume 2073 of LNCS, pages 772–784, San Francisco, CA, May 2001. Springer Verlag.